

UDC 316

Ulviyya Elyazova

Baku State University, PhD. student

ulviyyaali222@gmail.com

<https://orcid.org/0009-0007-4084-1057>

UNDERSTANDING EATING DISORDERS IS A COMPLEX MENTAL HEALTH CHALLENGE

The article discusses eating disorders in the modern era, a serious mental health issue related to weight and food. The most common types include anorexia nervosa and bulimia nervosa. These disorders can lead to severe physical and psychological consequences, including eating disturbances, bodily dysfunctions, and emotional distress. Anorexia nervosa is characterized by extreme food restriction and a fear of gaining weight. Bulimia nervosa involves episodes of excessive eating followed by purging. Eating disorders are influenced by genetic, psychological, social, and cultural factors. They are more common among adolescents and young adults, especially women, but can affect anyone differently. Early intervention is crucial for better treatment outcomes.

Purpose: The goal of understanding eating disorders is to recognize them as complex mental health issues that influence not only an individual's relationship with food, but also their emotional, psychological, and physical well-being.

Methodology: In the article studies the problem using the scientific method and comparative analysis methods.

Scientific novelty: The scientific novelty of understanding eating disorders as a complex mental health challenge stems from ongoing research that investigates the multifactorial nature of these conditions, integrating biological, psychological, and sociocultural factors in previously unknown ways.

Keywords: *anorexia, bulimia, mental health, social work, self-management, depression, anxiety*

Introduction

Eating disorders are serious mental health illnesses characterized by persistent changes in eating patterns, attitudes, and feelings around food, weight, and body image. These diseases can have serious psychological and physical effects, necessitating expert help to avoid long-term health problems. While eating disorders are typically associated with a fixation on weight and beauty, they are significantly more complicated and influenced by a range of biological, psychological, and social variables.

Anorexia nervosa is a well-known eating disorder that is distinguished by severe food restriction, a strong fear of gaining weight, and a skewed perspective of body image. Individuals with anorexia may perceive themselves as overweight, even if they are dangerously underweight. This condition can cause severe malnutrition, decreased immunological function, and, in extreme situations, organ failure. Similarly, bulimia nervosa is characterized by recurring episodes of binge eating, followed by compensatory behaviors such as self-induced vomiting, excessive exercise, or laxative abuse. Bulimia sufferers frequently experience a loss of control during binge episodes, as well as guilt and humiliation afterwards, which can lead to a deadly cycle of purging and restrictive behaviors.

Another common ailment is binge eating disorder (BED), which, unlike bulimia, does not include purging habits. Instead, people with BED eat a lot of food in a short period of time, sometimes even when they aren't hungry. This can result in obesity, diabetes, and other major health issues. In addition to these well-known diseases, Avoidant/Restrictive Food Intake

Disorder (ARFID) occurs when people avoid certain meals owing to sensory sensitivities, fear of choking, or a general lack of interest in eating. Unlike anorexia, ARFID is not motivated by body image concerns, but it nonetheless causes major nutritional shortages.

Eating disorders have multiple causes. Biological variables, such as genetic predisposition and neurotransmitter abnormalities, might raise a person's risk. Low self-esteem, perfectionism, anxiety, and depression are all key contributing causes. Furthermore, social and cultural forces have a role in the development of mental diseases, especially in civilizations that place an emphasis on excessive beauty standards. The pressure to obtain a specific body type, which is frequently reinforced by social media and popular culture, can lead to disordered eating practices in an attempt to meet these expectations. Furthermore, environmental factors including family relationships, childhood trauma, or a history of dieting can heighten sensitivity.

Eating disorders must be treated holistically and across all disciplines. Psychotherapy, particularly Cognitive Behavioral Therapy (CBT), has been demonstrated to help people identify and change negative thoughts about food and body image. Nutritional counseling is vital for reestablishing healthy eating habits and correcting nutritional deficits. In severe cases, medical attention may be required to manage issues such as heart disease, hormonal abnormalities, or organ damage. Furthermore, medication may be administered to treat co-occurring disorders such as depression or anxiety. Because eating disorders are frequently the result of emotional and psychological anguish, family, friends, and professionals must provide a supportive and compassionate environment for recovery.

Finally, eating disorders are more than just food or weight; they are complicated conditions that necessitate understanding, awareness, and professional intervention. Early diagnosis and intervention dramatically enhance recovery results, thus it is critical to educate people about the warning signs and encourage those who are struggling to seek assistance. We can help reduce the prevalence of eating disorders and support individuals in recovery by cultivating a society that values mental health, body acceptance, and adequate nutrition.

Anorexia Nervosa

Anorexia Nervosa is a severe eating disorder characterized by a strong fear of gaining weight, which causes people to severely restrict their food consumption. It is a complex disorder that affects both the mind and the body, frequently leading to major health problems [4, p. 189]. This illness primarily affects teens and young adults, particularly girls, however it can occur in persons of all ages and genders.

One of the hallmarks of Anorexia Nervosa is a skewed body image. People suffering with this illness frequently regard themselves as overweight, even if they are critically underweight. This view motivates individuals to engage in extreme behaviors such as excessive diets, fasting, and over-exercise. Some people may also use purging procedures such as vomiting or laxatives to avoid gaining weight. These practices are the result of fundamental psychological difficulties such as low self-esteem, perfectionism, and societal pressures that promote thinness.

Anorexia Nervosa has multiple causes, including genetic, environmental, and psychological aspects. According to studies, genetic predisposition may contribute to an increased risk of acquiring the condition. Furthermore, cultural factors that promote excessive beauty standards exacerbate body dissatisfaction, particularly among young people [6, p. 92]. Anxiety, sadness, and obsessive-compulsive tendencies all contribute to the illness, making it difficult for people to create a balanced connection with food and body image.

Anorexia Nervosa has potentially fatal consequences. Prolonged malnutrition causes serious physical issues such as heart disease, brittle bones, hormone abnormalities, and organ

failure. Many people endure acute exhaustion, dizziness, and irregular menstruation cycles because their bodies are unable to function effectively without appropriate nutrients. Furthermore, the psychological consequences are significant, typically resulting in social isolation, acute depression, and, in some cases, suicidal ideation.

Anorexia Nervosa is treated using a multidisciplinary approach that includes medical, psychological, and nutritional therapies. Patients in critical condition may require hospitalization to stabilize their health. Therapy, particularly cognitive-behavioral therapy (CBT), is widely utilized to assist people in addressing their skewed attitudes and habits linked to eating and body image. Nutritional counseling is essential for reinstating a healthy diet, and family therapy can help with rehabilitation.

Anorexia Nervosa is a complex and hazardous condition that requires timely diagnosis and treatment. It's more than just eating; it's a deep psychological struggle that has an impact on an individual's whole health. Raising awareness of the symptoms, causes, and implications of this condition is critical for preventing its onset and ensuring that those affected receive the appropriate care. Recovery is attainable with appropriate intervention and care, allowing people to regain both physical and emotional health.

Bulimia Nervosa is a complex eating disorder

Bulimia Nervosa is a severe eating disorder marked by episodes of binge eating followed by compensatory behaviors such as vomiting, excessive exercise, fasting, or laxative abuse. Individuals with this illness frequently lose control of their eating habits, resulting in episodes of excessive food consumption, which are subsequently counteracted by unhealthy techniques to prevent weight gain [5, p. 186]. Bulimia is a psychological illness that affects both the body and the psyche, often leading to serious health effects if not treated.

One of the distinguishing features of Bulimia Nervosa is an obsession with body weight and shape. People suffering from this illness may engage in hidden eating practices in response to emotions of shame and guilt connected with binge episodes. Unlike Anorexia Nervosa, Bulimia patients may maintain a normal weight or even be overweight, making it difficult to diagnose the illness based solely on looks. The binge-purge cycle is frequently fueled by mental turmoil, cultural constraints, and a crippling dread of gaining weight.

The causes of Bulimia Nervosa are complex, encompassing genetic, psychological, and environmental influences. A family history of eating problems, sadness, or anxiety increases the likelihood of developing Bulimia. Furthermore, cultural factors that promote thinness as the ideal body standard contribute to body dissatisfaction and disordered eating habits. Individuals with Bulimia are also more likely to experience psychological disorders such as low self-esteem, perfectionism, and impulsivity, which exacerbates their struggles with food and body image.

Bulimia Nervosa can have serious, even fatal, consequences. The regular cycle of bingeing and purging causes major medical consequences such as electrolyte imbalances, dehydration, gastrointestinal disorders, and teeth erosion due to stomach acid from continuous vomiting. Over time, these health issues might include cardiac abnormalities, kidney damage, and digestive problems. Furthermore, Bulimia has a major impact on mental health, raising the risk of anxiety, sadness, and substance dependence.

Treatment for Bulimia Nervosa necessitates a comprehensive strategy that treats both the medical and psychological elements of the condition [3, p. 77]. Cognitive-behavioral therapy (CBT) is commonly used to help people notice and improve their problematic eating habits, as well as their ideas about food and body image. Nutritional counseling is essential for developing a balanced diet, and drugs such as antidepressants may be administered to treat co-occurring mood problems. Individuals with Bulimia frequently experience feelings of isolation

and self-criticism, therefore support from family and friends is critical for rehabilitation.

To summarize, Bulimia Nervosa is a significant and complex eating disorder that requires prompt detection and treatment. It's more than just eating; it's a deeper psychological struggle that has an impact on an individual's overall health. Raising knowledge of its symptoms, causes, and repercussions is critical for giving support and treatment to people afflicted. With the right care, people can regain their physical and emotional health.

Binge-Eating Disorder (BED) is a severe and common eating disorder characterized by recurrent episodes of eating huge amounts of food in a short period of time, which are frequently accompanied by a loss of control and mental discomfort. Unlike Bulimia Nervosa, BED patients do not engage in compensatory behaviors such as purging or excessive exercise. This condition has serious psychological, physical, and societal ramifications, making it an urgent concern in modern healthcare.

The precise causes of BED are complex and varied, encompassing a variety of genetic, biochemical, psychological, and environmental variables. Genetics have a role in predisposing people to BED, with research showing that those with a family history of eating disorders or obesity are more likely to develop it. Furthermore, anomalies in brain function, particularly those related to impulse control and reward processing, contribute to disordered eating practices.

Depression, anxiety, low self-esteem, and prior trauma are all substantial risk factors for BED. Many people use food as a coping method when dealing with bad emotions or stress. Furthermore, societal pressures and cultural standards that encourage a negative relationship with food and body image might exacerbate the illness. Childhood experiences such as restrictive dieting, weight-related bullying, or parental disapproval may all raise the risk of developing BED.

Binge-Eating Disorder is diagnosed using specific criteria established in the Diagnostic and Statistical Manual of Mental Disorders (DSM-5). The key symptoms include frequent episodes of binge eating, which is defined as eating unusually large amounts of food in a short period of time, typically in secret. Individuals with BED frequently experience anxiety, humiliation, or guilt after binge-eating.

Other common symptoms include eating quickly, eating till uncomfortably full, eating when not hungry, and preferring to eat alone due to embarrassment. These activities must occur at least once a week for three months or longer to be classed as BED. Unlike other eating disorders, patients with BED do not engage in compensatory behaviors, which can lead to weight gain and accompanying health problems like obesity, diabetes, and cardiovascular disease.

BED has far-reaching repercussions for both physical and emotional health. Individuals with BED are more likely to be obese and have associated illnesses such as hypertension, high cholesterol, and metabolic disorders. Furthermore, the psychological consequences include an increased risk of mood disorders, social withdrawal, and lower quality of life. Individuals with BED frequently have low self-esteem and difficulties maintaining relationships as a result of the emotional load of their eating habits.

Effective therapy for BED requires a multidisciplinary strategy that addresses both the psychological and physiological aspects of the disorder [2, p. 268]. The most frequently suggested psychological treatment is Cognitive-Behavioral Therapy (CBT), which assists clients in recognizing and changing problematic food-related thought patterns and behaviors. Other therapeutic approaches, such as Interpersonal Therapy (IPT) and Dialectical Behavior Therapy (DBT), can help with underlying emotional disorders.

In some circumstances, medication is prescribed, such as selective serotonin reuptake inhibitors (SSRIs) and other appetite-regulating medicines. Nutritional counseling and lifestyle

changes, such as adopting healthier eating habits and participating in regular physical activity, are also important components of treatment. Support groups and peer counseling can offer emotional support and encouragement to those living with BED.

Binge-Eating Disorder is a complex and significant mental health issue that affects millions of people worldwide [1, p. 120]. Understanding its causes, symptoms, and effects is critical for timely discovery and intervention. Individuals suffering from BED can regain control of their eating habits and improve their overall health by combining therapy, medical treatment, and lifestyle modifications. Raising awareness and creating a supportive environment can help lessen the stigma associated with BED and enable those suffering from it to seek help without fear or shame.

Conclusion

To summarize, eating disorders are complicated mental health issues that extend beyond concerns about food and body image. They are characterized by deeply ingrained psychological, biological, and social elements that can have a major impact on a person's physical and emotional health. Anorexia nervosa, bulimia nervosa, binge eating disorder, and avoidant/restrictive food intake disorder (ARFID) all present unique issues, but they are linked by distress and inappropriate eating practices. Understanding the varied nature of eating disorders is critical for their prevention and treatment.

A multidisciplinary strategy is required for effective treatment, which may include psychotherapy, nutritional counseling, medical monitoring, and, on occasion, medication. It is also critical to realize the value of early intervention, as it can significantly enhance recovery rates and prevent long-term health consequences. We can help people suffering from eating disorders obtain the help they need and improve their quality of life by cultivating a society that values mental health awareness, body acceptance, and a balanced attitude to food. Finally, eating disorders are not just about food; they are indications of deeper emotional and psychological issues that necessitate compassionate and comprehensive treatment.

References

1. "Eating Disorders: The Facts" by Helen M. V. King (2016), 160 pages
2. "Intuitive Eating: A Revolutionary Program That Works" (4th edition) by Evelyn Tribole and Elyse Resch (2020), 368 pages
3. "Just Eat It: How Intuitive Eating Can Help You Get Together Around Food" by Laura Thomas (2018), 336 pages
4. "Radical Recovery: A Political Guide to Healing from Eating Disorders" by Stephanie Covington Armstrong (2017), 300 pages
5. "The Body Image Workbook: An 8-Step Program for Learning to Like Your Looks" by Thomas F. Cash (2018), 296 pages
6. "The Psychology of Eating: From Healthy to Disordered Behavior" by Jane Ogden (2016), 288 pages

Ülviyyə Elyazova

QIDA QƏBULU POZUNTULARINI ANLAMAQ KOMPLEKS ƏQLİ SAĞLAMLIQ PROBLEMİ KİMİ

Xülasə

Məqalədə müasir dövrdə qida qəbul etmək pozuntusundan, çəki və qida ilə bağlı ciddi əqli sağlamlıq problemlərindən bəhs edilir. Ən çox rast gəlinən növlərə anoreksiya nervoza və bulimiya nervoza daxildir. Bu pozuntular qidalanma, orqanizmdə yaranan problemlərlə və emosional narahatlıq da daxil olmaqla ciddi fiziki və psixoloji nəticələrə səbəb ola bilər. Anoreksiya nervoza həddindən artıq qida məhdudiyyəti, kökəlmək qorxusu ilə xarakterizə olunur. Bulimiya nervoza, həddindən artıq yemək və qaytarma ilə bağlıdır. Yemək pozuntusuna genetik, psixoloji, sosial və mədəni amillər təsir edir. Onlar yeniyetmələrdə və gənclərdə, xüsusən də qadınlarda daha çox rast gəlinir, lakin hər kəsə fərqli təsir edə bilər. Erkən müdaxilə daha yaxşı müalicə nəticələri üçün çox vacibdir.

Məqsəd: Məqalədə yemək pozuntusunu başa düşməkdə məqsəd onları tək cə insanın qida ilə münasibətinə deyil, həm də onların emosional, psixoloji və fiziki rifahına təsir edən kompleks psixi sağlamlıq problemləri kimi tanımaqdır.

Metodologiya: Məqalədə elmi metod və müqayisəli təhlil metodları ilə problem tədqiq olunur.

Elmi yenilik: Məqalədə qida pozuntularının kompleks psixi sağlamlıq problemi kimi başa düşülməsinin elmi yeniliyi bu şərtlərin multifaktorial təbiətini araşdıran, bioloji, psixoloji və sosial-mədəni amilləri əvvəllər məlum olmayan üsullarla birləşdirən davamlı tədqiqatlardan qaynaqlanır.

Açar sözlər: anoreksiya, bulimiya, psixi sağlamlıq, sosial iş, özünüidarəetmə, depressiya, təşviş

Ульвия Эльязова

ПОНИМАНИЕ РАССТРОЙСТВ ПИЩЕВОГО ПОВЕДЕНИЯ-СЛОЖНАЯ ПРОБЛЕМА ПСИХИЧЕСКОГО ЗДОРОВЬЯ

Резюме

В статье рассматриваются расстройства пищевого поведения в наше время — серьезная проблема психического здоровья, связанная с весом и питанием. Наиболее распространёнными типами являются нервная анорексия и нервная булимия. Эти расстройства могут привести к серьёзным физическим и психологическим последствиям, включая проблемы с питанием, заболевания внутренних органов и эмоциональный стресс. Нервная анорексия характеризуется крайним ограничением в еде и страхом набрать вес. Нервная булимия связана с перееданием и очищением желудка. На расстройства пищевого поведения влияют генетические, психологические, социальные и культурные факторы. Они чаще встречаются у подростков и молодых людей, особенно у женщин, но могут проявляться у каждого по-разному. Раннее вмешательство имеет решающее значение для достижения лучших результатов лечения.

Цель: Целью понимания расстройств пищевого поведения является признание их как сложных проблем психического здоровья, которые влияют не только на отношение человека к еде, но и на его эмоциональное, психологическое и физическое благополучие.

Методология: В статье проблема изучается с использованием научного метода и методов сравнительного анализа.

Научная новизна: Научная новизна понимания расстройств пищевого поведения как сложной проблемы психического здоровья вытекает из продолжающихся исследований, которые изучают многофакторную природу этих состояний, объединяя биологические, психологические и социокультурные факторы ранее неизвестными способами.

Ключевые слова: анорексия, булимия, психическое здоровье, социальная работа, самоменеджмент, депрессия, тревожность